

## INFRARED LASER SPECTROSCOPY OF THE n-PROPYL AND i-PROPYL RADICALS IN HELIUM DROPLETS: SIGNIFICANT BEND-STRETCH COUPLING REVEALED IN THE CH STRETCH REGION

CHRISTOPHER P. MORADI, GARY E. DOUBERLY, *Department of Chemistry, University of Georgia, Athens, GA, USA*; DANIEL P. TABOR, EDWIN SIBERT, *Department of Chemistry, The University of Wisconsin, Madison, WI, USA*.

The n-propyl and i-propyl radicals were generated in the gas phase via pyrolysis of n-butyl nitrite ( $\text{CH}_3(\text{CH}_2)_3\text{ONO}$ ) and i-butyl nitrite ( $\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{ONO}$ ) precursors, respectively. Nascent radicals were promptly solvated by a beam of He nanodroplets, and the infrared spectra of the radicals were recorded in the C-H stretching region. In addition to three vibrations of n-propyl previously measured in an Ar matrix, we observe many unreported bands between 2800 and 3150  $\text{cm}^{-1}$ , which we attribute to propyl radicals. The C-H stretching modes observed above 2960  $\text{cm}^{-1}$  for both radicals are in excellent agreement with anharmonic frequencies computed using VPT2. Between 2800 and 2960  $\text{cm}^{-1}$ , however, the spectra of n-propyl and i-propyl radicals become quite congested and difficult to assign due to the presence of multiple anharmonic resonances. Computations employing a local mode Hamiltonian reveal the origin of the spectral congestion to be strong coupling between the high frequency C-H stretching modes and the lower frequency bending/scissoring motions. The only significant local coupling is between stretches and bends on the same  $\text{CH}_2/\text{CH}_3$  group.